



by Allard USA




## ARM SLING with THUMB LOOP

Rx Only — Single Patient Use Only



### Instructions for Use



**FEEL THE FUNCTION**

---

*The intended function of this sling is to transfer the weight of the forearm to the patient's back and unaffected shoulder. Review instructions on back page carefully before application.*

---



by Allard USA




## ARM SLING with THUMB LOOP

Rx Only — Single Patient Use Only



### Instructions for Use



**FEEL THE FUNCTION**



---

*The intended function of this sling is to transfer the weight of the forearm to the patient's back and unaffected shoulder. Review instructions on back page carefully before application.*


---

## Arm Sling with Thumb Loop

1. Slip strap end through both D-rings and secure loosely.



2. Place affected arm into sling.  
Loop strap over head.



3. Make sure elbow is seated well into corner pocket of sling. Slide thumb into thumb loop. Adjust strap to hold forearm/wrist at angle recommended by clinician.

### Indications / Intended Use

Cast support; Injuries to the arm, wrist or hand; Acromioclavicular sprains; Anterior glenohumeral dislocations.

### Contraindications

Inability to bend arm to at least 90°.

### Wash instruction

Hand wash in cold water using mild detergent.  
Air dry.

### Adjustment

This sling should only be fitted by a licensed medical professional.


### Warning

If wearing the sling causes any unusual reactions, swelling or prolonged pain, remove and consult your medical professional. Check skin on a daily basis.


**CSUS**  
by Allard USA

## Arm Sling with Thumb Loop

1. Slip strap end through both D-rings and secure loosely.



2. Place affected arm into sling.  
Loop strap over head.



3. Make sure elbow is seated well into corner pocket of sling. Slide thumb into thumb loop. Adjust strap to hold forearm/wrist at angle recommended by clinician.

### Indications / Intended Use

Cast support; Injuries to the arm, wrist or hand; Acromioclavicular sprains; Anterior glenohumeral dislocations.

### Contraindications

Inability to bend arm to at least 90°.

### Wash instruction

Hand wash in cold water using mild detergent.  
Air dry.

### Adjustment

This sling should only be fitted by a licensed medical professional.

### Warning

If wearing the sling causes any unusual reactions, swelling or prolonged pain, remove and consult your medical professional. Check skin on a daily basis.

**CSUS**  
by Allard USA