

Hyperbaric Oxygen Therapy for Post-Concussion Syndrome: Contradictory Conclusions from a Study Mischaracterized as Sham-Controlled

Paul G. Harch

Dear Editor,

The recent study by Wolf and associates¹ has affirmed the effectiveness of hyperbaric (oxygen) therapy in the treatment of patients with mild traumatic brain injury (mTBI)/post-concussion syndrome (PCS) and post-traumatic stress disorder (PTSD). This affirmation emerges from analysis of the study data, rather than from the study's stated conclusions. Mischaracterized as a sham-controlled (placebo implied) design, the study errs in concluding that "HBO₂ at 2.4 ATA pressure had no effect on post-concussive symptoms after mild TBI." A reconsideration of the science of hyperbaric therapy reveals that the study by Wolf and colleagues¹ is neither a sham nor placebo-controlled study. Rather, it is a Phase II study of two composite doses of hyperbaric therapy that demonstrated significant improvements in PCS and PTSD symptoms at the 2.4 atmospheres absolute (ATA) pure oxygen dose as well as the low-pressure 1.3 ATA air/oxygen dose.

Hyperbaric (oxygen) therapy (HBOT) is a combination product of increased pressure and increased pressure of oxygen above ambient atmospheric pressure, according to scientific principles and current Food and Drug Administration understanding. Although traditionally misdefined as a treatment for diseases based on the increased oxygen component alone (> 1.4 ATA oxygen),² it is a treatment with hyperbaric pressure and hyperoxia for disease processes^{2,3} whose primary targets are oxygen and pressure sensitive genes.^{4–6}

Evidence for this dual component nature of hyperbaric therapy is found in the 351-year history of hyperbaric air therapy⁷ and the recent 60-year history of animal, human tissue, and human experiments that have documented biological effects of pressure, especially in the micropressure range^{8,9} of the Wolf and coworkers¹ "sham" control group and the control groups of the Department of Defense (DoD) HBOT TBI studies.¹⁰ Examples of this literature are listed in Tables 1 and 2.^{11–28} Pressures from 1.21–1.26 ATA delivered to human^{29–31} and 1.0015–1.015 ATA to animal endothelial cells,³² and 1.10 and 1.20 ATA to human platelets^{33,34} for 15 min or longer have caused the elaboration or suppression of vasoactive substances,^{29–31} and the elaboration of growth factors,³² inflammatory mediators,³³ oxidation products,³⁴ and cell proliferation.³² This literature and biological effects from a 1-min exposure

to 1.09 ATA or 3 min at 1.04 ATA¹⁷ inform the symptomatic improvements noted in the Wolf and associates¹ "sham" group, as do benefits of hyperbaric air on spinal function and PTSD in spinal cord injured veterans during a SCUBA diving training course.²⁸

To meet the definition of a true sham,³⁵ any controlled experiment to test HBOT must omit in its control groups the active ingredients of increased pressure and hyperoxia. The Wolf and colleagues¹ "sham" control group does neither; rather, it includes both. The "sham" control group is exposed to 1.3/1.2 ATA of air, which is a 20–30% increase in pressure and 28–43% increase in plasma oxygen³⁶ over sea level plasma oxygen and a slightly greater increase over San Antonio (hyperbaric treatment site) atmospheric pressure.³⁷ Because pressure and hyperoxia are non-inert—i.e., are biologically active—the Wolf and coworkers¹ "sham" control group cannot test for placebo effects; placebo/placebo response is defined as "The effect that an inactive or inert substance has on a clinical condition."³⁸ Wolf and associates¹ allude to possible bioactivity of the control group, but the lack of discussion indicates a lack of appreciation that the presence of hyperoxia and pressure negate Wolf and colleagues¹ characterization as a "sham" control group.

Restating the design of the Wolf and coworkers¹ study, it is a Phase II comparative dosing study of two composite doses of hyperbaric therapy (four actual doses), compressed air (low dose increased pressure and increased oxygen), and compressed oxygen (high dose pressure and high dose oxygen). Both doses were efficacious in the treatment of mTBI PCS and PTSD. The PTSD data demonstrated 18% and 22% reductions in the PCL-M (interpolated from the Figure 1 graph in Wolf and colleagues¹) in the HBOT and "sham" groups, respectively, after 30 2 h treatments. These reductions compared favorably with five other therapies/six studies for PTSD^{39–44} that used the PCL-M (6–45% reductions).

The PCS ImPACT data were similarly significantly improved in both groups, but it is the disparity in component and pattern change on the ImPACT results for the two groups that underscore the dual dose design of the study and efficacy of these two doses: 10 ImPACT scores significantly improved in the low dose group compared with 2 in the high dose group. For all 22 items of the ImPACT, 20 improved, 1 was unchanged, and 1 was worse in the

TABLE 1. IN VITRO AND IN VIVO STUDIES ON HYPERBARIC (PRESSURE AND OXYGEN) EFFECTS

<i>Study</i>	<i>Year</i>	<i>Model: In Vivo (IV), In Vitro (IVT)</i>	<i>Animal/human</i>	<i>Ambient gas</i>
1. Cunningham, O.J. ¹¹	1900–1929	Multiple medical disorders	Human	Air
2. Dowell, R.T. ¹²	1978	Aortic constriction (IV)	Rat-neonate and adult	Air
3. Hishikawa, K. ¹³	1994	Vascular smooth muscle cells, IVT	Rat	Helium
4. Mattana, J. ¹⁴	1995	Mesangial (renal) cells (IVT)	Rat	95% air, 5% CO ₂
5. Teiger, E. ¹⁵	1996	Aortic constriction (IV)	Rat	Air
6. Kramer, M.R. ¹⁶	1998	COPD-pulmonary function	Humans	Air
7. Kawata, Y. ¹⁷	1998	Mesangial (renal) cells (IVT)	Rat	Air
8. Macdonald, A.G.-review ⁹	1999	Wide range of IV and IVT	Bacteria, insect, nematode, animal, human	95% air, 100% air, helium
9. Agar, A. ¹⁸	2000	Neuroblastoma (IVT), glaucoma model	Rat	95% air, 5% CO ₂
10. Dean, J.B. ¹⁹	2000	Brain stem neurons (IVT)	Rat	Air, helium
11. Heal, R.D. ²⁰	2001	Dorsal root ganglion neurons (IVT)	Rat	95% air, 5% CO ₂
12. Collet, J.P. ²¹	2001	Cerebral palsy (IV)	Human-children	Air, 100% oxygen
13. Yu, H.A.I. ²²	2002	Subacute hypoxic-ischemia brain injury (IV)	Rat	Air, 100% oxygen
14. Heuser, G. ²³	2002	Chronic toxic encephalopathy or autism (IV)	Human-adults and children	24% oxygen
15. Kazantseva, N.V. ²⁴	2002	Cerebrovascular injury, epilepsy, migraine (IV)	Human adults	30%, 40%, and 100% oxygen
16. Rusyniak, D.E. ²⁵	2003	Acute stroke	Human-adults	100% oxygen
17. Stanley, A.C. ²⁶	2005	Fibroblast wound-healing (IVT)	Human neonatal fibroblasts	94% air, 6% CO ₂
18. Rossignol, D.A. ²⁷	2009	Autism	Human-children	Air, 24% oxygen
19. Kaplin, A. ²⁸	2011	Spinal cord injury and PTSD	Human-veterans	Air

COPD, chronic obstructive pulmonary disease; PTSD, post-traumatic stress disorder.

low dose group, while 11 improved, 3 were unchanged, and 8 were worse in the high dose group (Table 1 in Wolf and associates¹). The pattern of composite ImPACT scores over the course of the study is also different for the two groups (Figure 2 in Wolf and coworkers¹). Subjects in the low dose group experienced initial deterioration then steady improvement until the end of the study while the high dose group showed improvement then a steady reversal of benefit to near baseline, followed by rebound improvement 6 weeks post-treatment.

This sinusoidal trajectory in the high dose group suggests a differential dosing effect and possibly an overdose response with partial recovery after removal of the high dose, and is consistent with the worsening of eight scores in this group. It is also consistent with a phenomenon previously described and documented in multiple cases of HBOT treatment of chronic cerebral disorders.⁴⁵

Since the initial submission of this Letter to the Editor, Wolf and colleagues¹ have now confirmed this overdosing effect on PCS symptoms in the 2.4 ATA group in a subset analysis presented on June 14, 2013.⁴⁶ This demonstration of low dose effectiveness and progressive high dose overdosing is also evident in other animal and clinical studies.^{25,47,48}

Wolf and associates¹ list multiple possibilities for the improvements in PCS and PTSD, including “placebo, Hawthorne effect, the natural resolution of symptoms over time,...exposure to sham-control partial pressures of oxygen and nitrogen,...and change in living environment...and daily routine.” While 30 chamber experiences and change in living environment and daily routine are theoretical placebo contributions, the myriad standard clinical pressure/hyperoxia² and micropressure-induced biological effects demonstrated in both animal and human studies^{8,9} and Tables 1 and 2 suggest that placebo is not the majority effect.

At the same time, placebo effects were also present in all six of the aforementioned PTSD/PCL-M studies. Hawthorne effect is also

theoretically possible, but the foundation/proving of the Hawthorne effect has been undermined by re-evaluation of the initial data set.⁴⁹ Symptoms in veterans with TBI and PTSD, in fact, do not resolve over time, as acknowledged by Wolf and coworkers¹ in their conclusion “improved more than would be expected greater than 6 months after mTBI,” and reports on persistent care and even worsening of condition over time in the Veterans Affairs system.^{50,51}

All of the explanations by Wolf and colleagues,¹ however, do not explain the disparity in component and pattern change in the ImPACT data and transient overdose effect in the high dose oxygen group. If placebo, Hawthorne effect, and other non-biological etiologies are causally entertained, why wouldn't the component and pattern changes be identical in the two groups? The last choice, effects of increased partial pressures of oxygen and nitrogen, seems most plausible as the dominant etiology for the statistically significant improvements in both groups, but it is not necessarily nitrogen pressure, but pressure, per se, that is involved. Wolf and associates¹ state, however, “...it seems very unlikely such a minimal dose of oxygen and nitrogen could influence brain function favorably.” The preceding argument and studies in Table 1 and 2 suggest the opposite.

In conclusion, the study by Wolf and colleagues¹ is a non-sham/non-placebo/non-controlled Phase II two composite dose study of hyperbaric therapy (hyperbaric air and hyperbaric oxygen) in U.S. veterans with PCS from mTBI with or without PTSD. The study demonstrated significant net improvements in PCS and PTSD symptoms with both doses of hyperbaric therapy, improvements that are similar in magnitude to other therapies for PTSD and greater than would be expected for PTSD and PCS over time without treatment based on the natural history of the diseases and published persistence rates in veterans.^{50,51} Their results are thus comparatively effective to other existing therapies for PTSD and possibly PCS of mTBI.

TABLE 2. CONTINUATION OF TABLE 1 IN VITRO AND IN VIVO STUDIES ON HYPERBARIC (PRESSURE AND OXYGEN) EFFECTS

Study	Oxygen pressure (ATA): Control:experimental	Total pressure (ATA): Control:experimental	Duration/# Rxs	Effect
1.	No controls:0.281-0.496	No controls: average1.34-2.36 ATA, max 3 ATA	2-3 hours up to days. Multiple treatments	Generally, improvement in clinical and laboratory parameters
2.	0.129:0.129 (arterial oxygen partial pressure)	1.103-1.163: 1.150-1.217	2,3,4,5 weeks	Cardiac myocyte and non-myocyte proliferation
3.	0.205:0.205	1.000:1.053, 1.105, 1.158	24 and 48 h	Increased cell proliferation and DNA synthesis; process begins w/i 30s press.
4.	0.23:0.237, not measured for 2nd pressure	1.000:1.053-1.066 and 1.066-1.079	1 or 7 days	Decreased cell number and increased matrix collagen (MC) at 7d. Dose-response for MC.
5.	0.129:0.129 (arterial oxygen partial pressure)	Approximately same as #1; same model	1,2,4,7,15,30 days	Increased apoptosis of cardiac myocytes by 1 day, peak at 4 days
6.	No controls (pre/post):0.221	Pre: 0.912 Post: 1.0526	21 days	Improved arterial oxygen and exercise capacity.
7.	0.270:0.270	1.000:1.039, 1.066, 1.092, 1.118	1-30 minutes, 12, 24, 48 hours	Enzyme activation w/i 1 min @ 1.092 ATA, gene expression/ proliferation post 12 h and 24 h exposure
8.	0.199:0.251	1.000:1.010-1.260	Few sec-days	Wide range of biological effects
9.	0.222:0.222	1.000:1.131	2h	Increased apoptosis
10.	0.950:2.352-3.332	1.000:1.000-4.000	5-15 min	Differential effects of pressure and oxygen on firing rate
11.	0.199:0.599	1.000:3.000	8 min 20 sec	Suppression of firing rate
12.	0.273:1.750	1.300:1.750	60 minutes/dx40 d	Both groups: Significant improvements in motor and cognition
13.	0.250:0.625, 2.50	1.000:2.500	60 min/day x 10 days	Increased bFGF in hyperbaric air, increased bFGF and bFGF mRNA in HBO
14.	No control: 0.312	No control: 1.30	1 h/day x 10 days	Improved blood flow on SPECT brain imaging
15.	0.300:0.330, 0.480, 1.500	1.000:1.100, 1.200, 1.500	15-20m? x 5: 15-20m x 4-10, 20 m x 1 or 4-5, 60 m x 5	Improvement symptoms, EEG, blood flow, vital signs, biochemical params. < 1.5 ATA
16.	1.14:2.50	1.14:2.50	1 h	Better outcome in the low dose group
17.	0.194:0.205, 0.213, 0.229	1.000:1.039, 1.079, 1.158	12-18 h	Delayed wound healing and less proliferating cell nuclear antigen
18.	0.216: 0.312	1.03:1.30	1 h/day x 40 days	Clinical improvement
19.	SCUBA course	Likely < 3 ATA	Likely < 1 h	Improved spinal cord function and PTSD

ATA, atmospheres absolute; bFGF, basic fibroblast growth factor; HBO, hyperbaric oxygen; EEG, electroencephalography; PTSD, post-traumatic stress disorder.

If further evidence for the efficacy of hyperbaric therapy in mTBI PCS and PTSD is deemed necessary, as alternatives to the “pivotal” trial,¹⁰ and in light of nearly \$1 billion dollars of research and development money already spent by the Department of Defense (DoD) on research seeking effective treatments for TBI and PTSD, this author would suggest two options: (1) Abandoning the pursuit of a “sham” hyperbaric control group and prioritizing the data from the wait-list group in the DoD HOPP study¹⁰ (Wolf and coworkers¹ mention a wait-list group in their recommendations for future studies). This option is based on the physical impossibility of controlling for a chamber experience because of the inability to control for pressure (no method to duplicate middle ear pressure changes without placement of pressure equalization tubes and no commercially available mechanisms to duplicate adiabatic heating and cooling on compression and decompression). Continued attempts at a “sham” pressure control group will only further confuse the scientific and lay community; and (2) an economical Civilian/DoD/Veterans Affairs (VA) off-label networked hyperbaric treatment program using a Medicare-like Coverage with Evidence Development⁵² pathway. The safety of HBOT 1.5 and even higher doses is not in issue; hyperbaric oxygen therapy in mTBI PCS/PTSD has satisfied one of the cardinal rules of medicine, “First, Do No Harm.” The Coverage with Evidence pathway would allow the DoD and VA to immediately begin treating active military and veteran casualties with hyperbaric therapy in both military and civilian clinics/hospitals such that this health and quality of life improving therapy can be delivered without further delay to the hundreds of thousands of injured PCS/PTSD casualties and veterans now in need.

Author Disclosure Statement

Dr. Harch owns a small consulting company, Harch Hyperbarics, Inc., and is president of a non-profit corporation, The International Hyperbaric Medical Foundation (IHMF). He derives no income, salary, or benefits from the IHMF.

References

- Wolf, G., Cifu, D., Baugh, L., Carne, W., and Profenna, L. (2012). The effect of hyperbaric oxygen on symptoms after mild traumatic brain injury. *J. Neurotrauma* 29, 2606–2612.
- Gesell, L.B., ed. (2009). *Hyperbaric Oxygen Therapy Indications*, 12th ed. The Hyperbaric Oxygen Therapy Committee Committee Report. Undersea and Hyperbaric Medical Society: Durham, NC, pps. 4–5.
- Harch, P.G., and Neubauer, R.A. (1999). Hyperbaric oxygen therapy in global cerebral ischemia/anoxia and coma, in: Chapter 18, *Textbook of Hyperbaric Medicine*, 3rd revised ed. K.K. Jain (ed.). Hogrefe and Huber Publishers: Seattle, pps. 319–345.
- Godman, C.A., Chheda, K.P., Hightower, L.E., Perdrietz, G., Shin, D.G., and Giardina, C. (2010). Hyperbaric oxygen induces a cytoprotective and angiogenic response in human microvascular endothelial cells. *Cell Stress Chaperones* 15, 431–442.
- Chen, Y., Nadi, N.S., Chavko, M., Auker, C.R., and McCarron, R.M. (2009). Microarray analysis of gene expression in rat cortical neurons exposed to hyperbaric air and oxygen. *Neurochem. Res.* 34, 1047–1056.
- Kendall, A.C., Whatmore, J.L., Harries, L.W., Winyard, P.G., Eggleston, P., and Smerdon, G.R. (2013). Different oxygen treatment pressures alter inflammatory gene expression in human endothelial cells. *Undersea Hyper. Med.* 40:115–123.
- Jain, K.K. (2004). The history of hyperbaric medicine, in: *Textbook of Hyperbaric Medicine*, 4th revised and expanded edition, K.K. Jain (ed.). Hogrefe and Huber Publishers: Cambridge, MA, pps. 3–8.
- Dean, J.B., Mulkey, D.K., Garcia, A.J., III, Putnam, R.W., and Henderson, R.A., III. (2003). Neuronal sensitivity to hyperoxia, hypercapnia, and inert gases at hyperbaric pressures. *J. Appl. Physiol.* 95, 883–909.
- Macdonald, A.G., and Fraser, P.J. (1999). The transduction of very small hydrostatic pressures. *Comp. Biochem. Physiol. A. Mol. Integr. Physiol.* 122, 13–36.
- Weaver, L.K., Cifu, D., Hart, B., Wolf, G., and Miller, S. (2012). Hyperbaric oxygen for post-concussion syndrome: design of Department of Defense clinical trials. *Undersea Hyperb. Med.* 39, 807–814.
- Trimble, V.H. (1974). *The Uncertain Miracle: Hyperbaric Oxygenation*. Doubleday & Company, Inc.: Garden City, NY.
- Dowell, R.T., and McManus, R.E., III. (1978). Pressure-induced cardiac enlargement in neonatal and adult rats. Left ventricular functional characteristics and evidence of cardiac muscle cell proliferation in the neonate. *Circ. Res.* 42, 303–310.
- Hishikawa, K., Nakaki, T., Marumo, T., Hayashi, M., Suzuki, H., Kato, R., and Saruta, T. (1994). Pressure promotes DNA synthesis in rat cultured vascular smooth muscle cells. *J. Clin. Invest.* 93, 1975–1980.
- Mattana, J., and Singhal, P.C. (1995). Applied pressure modulates mesangial cell proliferation and matrix synthesis. *Am. J. Hypertens.* 8, 1112–1120.
- Teiger, E., Than, V.D., Richard, L., Wisniewsky, C., Tea, B.S., Gaboury, L., Tremblay, J., Schwartz, K., and Hamet, P. (1996). Apoptosis in pressure overload-induced heart hypertrophy in the rat. *J. Clin. Invest.* 97, 2891–2897.
- Kramer, M.R., Springer, C., Berkman, N., Glazer, M., Bublil, M., Bar-Yishay, E., and Godfrey, S. (1998). Rehabilitation of hypoxic patients with COPD at low altitude at the Dead Sea, the lowest place on earth. *Chest* 113, 571–575.
- Kawata, Y., Mizukami, Y., Fujii, Z., Sakumura, T., Yoshida, K., and Matsuzaki, M. (1998). Applied pressure enhances cell proliferation through mitogen-activated protein kinase activation in mesangial cells. *J. Biol. Chem.* 273, 16905–16912.
- Agar, A., Yip, S.S., Hill, M.A., and Coroneo, M.T. (2000). Pressure related apoptosis in neuronal cell lines. *J. Neurosci. Res.* 60, 495–503.
- Dean, J.B., and Mulkey, D.K. (2000). Continuous intracellular recording from mammalian neurons exposed to hyperbaric helium, oxygen, or air. *J. Appl. Physiol.* 89, 807–822.
- Heal, R.D., Rogers, A.T., Lunt, G.G., Pointer, S.A., and Parsons, A.T. (2001). Development of a neuronal pressure sensor. *Biosens. Bioelectron.* 16, 905–909.
- Collet, J.P., Vanasse, M., Marois, P., Amar, M., Goldberg, J., Lambert, J., Lassonde, M., Hardy, P., Fortin, J., Tremblay, S.D., Montgomery, D., Lacroix, J., Robinson, A., and Majnemer, A., HBO-CP Research Group. (2001). Hyperbaric oxygen for children with cerebral palsy: a randomized multicentre trial. *Lancet* 357, 582–586.
- Yu, H.A.I., Tian, R.L., Pan, X.W., Luan, Z., and Song, L.W. (2002). Effects of hyperbaric oxygen on brain bFGF and bFGF mRNA expression of neonatal rats after hypoxia-ischemia injury, in: *The Proceedings of the 2nd International Symposium on Hyperbaric Oxygenation for Cerebral Palsy and the Brain-Injured Child*. J.T. Joiner (ed). Best Publishing Co: Flagstaff, pps. 57–66.
- Heuser, G., Heuser, S.A., Rodlander, D., Aguilera, O., and Uszler, M. (2002). Treatment of neurologically impaired adults and children with “mild” hyperbaric oxygen (1.3ATA and 24% oxygen), in: *The Proceedings of the 2nd International Symposium on Hyperbaric Oxygenation for Cerebral Palsy and the Brain-Injured Child*. J.T. Joiner (ed). Best Publishing Co: Flagstaff, pps. 109–116.
- Kazantseva, N.V. (2002). Mechanisms of curative effect of minimized hyperbaric treatment in cerebral ischemia, in: *The Proceedings of the 2nd International Symposium on Hyperbaric Oxygenation for Cerebral Palsy and the Brain-Injured Child*. J.T. Joiner (ed). Best Publishing Co: Flagstaff, pps. 199–212.
- Rusyniak, D.E., Kirk, M.A., May, J.D., Kao, L.W., Brizendine, E.J., Welch, J.L., Cordell, W.H., and Alonso, R.J. (2003). Hyperbaric oxygen therapy in acute ischemic stroke: results of the Hyperbaric Oxygen in Acute Ischemic Stroke Trial Pilot Study. *Stroke* 34, 571–574.
- Stanley, A.C., Lounsbury, K.M., Corrow, K., Callas, P.W., Zhar, R., Howe, A.K. and Ricci, M.A. (2005). Pressure elevation slows the fibroblast response to wound healing. *J. Vasc. Surg.* 42, 546–551.
- Rosignol, D.A., Rosignol, L.W., Smith, S., Schneider, C., Logerquist, S., Usman, A., Neubrandner, J., Madren, E.M., Hintz, G., Grushkin, B., and Mumper, E.A. (2009). Hyperbaric treatment for children with autism: a multicenter, randomized, double-blind, controlled trial. *BMC Pediatr.* 13, 9–21.

28. Kaplin, A., and Becker, D. Scuba diving improves function of body, mind in vets with spinal cord injury. Available at: <http://www.newswise.com/articles/view/580607>. Accessed September 21, 2013.
29. Tokunaga, O., and Watanabe, T. (1987). Properties of endothelial cells and smooth muscle cell cultured in ambient pressure. *In Vitro Cell Dev. Biol.* 23, 528–534.
30. Tokunaga, O., Fan, J.L., and Watanabe, T. (1989). Atherosclerosis and endothelium. Part II. Properties of aortic endothelial and smooth muscle cells cultured at venous ambient pressures. *Acta Pathol Jpn*, 39, 356–362.
31. Hishikawa, K., Nakaki, T., Suzuki, H., Saruta, T., and Kato, R. (1992). New method of investigating functional roles of pressure sensitive mechanoreceptor in human endothelial cells. *J. Cardiovasc. Pharmacol.* 20, Suppl. 12, S66–S67.
32. Acevedo, A.D., Bowser, S.S., Gerritsen, M.E., and Bizios, R. (1993). Morphological and proliferative responses of endothelial cells to hydrostatic pressure: role of fibroblast growth factor. *J. Cell Physiol.* 157, 603–614.
33. Torsellini, A., Belvicci, A., Citti, S., Cozzolino, F., Guidi, G., Lombardi, V., Vercilli, D., and Veloci, M. (1982). Effects of pressure excursions on human platelets. In vitro studies on beta-thromboglobulin (β -TG) and platelet factor 4 (PF4) release and on platelet sensitivity to ADP-aggregation. *Haematologica* 67, 860–866.
34. Torsellini, A., Donc, L., Palinski, W., Guidi, G., and Lombardi, V. (1984). Pressure-induced metabolic activation of platelets demonstrated in vitro by determination of malondialdehyde formation. *Thromb. Haemostas.* 52, 4–6.
35. Sham. Available at: <http://www.merriam-webster.com/medical/sham>. Accessed September 13, 2013.
36. Alveolar gas equation. Available at: http://en.wikipedia.org/wiki/Alveolar_gas_equation. Accessed September 13, 2013.
37. San Antonio. Available at: http://en.wikipedia.org/wiki/San_Antonio#Geography. Accessed September 21, 2013.
38. Placebo effect. Available at: <http://medical-dictionary.thefreedictionary.com/placebo+effect>. Accessed September 21, 2013.
39. Alvarez, J., Mclean, C., Harris, A.H., Rosen, C., Ruzek, J.I., and Kimerling, R. (2011). The comparative effectiveness of cognitive processing therapy for male veterans treated in a VHA posttraumatic stress disorder residential rehabilitation program. *J. Consult. Clin. Psychol.* 79, 590–599.
40. Chard, K.M., Schumm, J.A., McIlvain, S.M., Bailey, G.W., and Parkinson, R.B. (2011). Exploring the efficacy of a residential treatment program incorporating cognitive processing therapy-cognitive for veterans with PTSD and traumatic brain injury. *J. Trauma. Stress* 24, 347–351.
41. Macdonald, A., Monson, C.M., Doron-Lamarca, S., Resick, P.A., and Palfai, T.P. (2011). Identifying patterns of symptom change during a randomized controlled trial of cognitive processing therapy for military-related posttraumatic stress disorder. *J. Trauma. Stress* 24, 268–276.
42. Rizzo, A., Parsons, T.D., Lange, B., Kenny, P., Buckwalter, J.G., Rothbaum, B., Difede, J., Frazier, J., Newman, B., Williams, J., and Reger, G. (2011). Virtual reality goes to war: a brief review of the future of military behavioral healthcare. *J. Clin. Psychol. Med. Settings* 18, 176–187.
43. Rosenthal, J.A., Grosswald, S., Ross, R., and Rosenthal, N. (2011). Effects of transcendental meditation in veterans of Operation Enduring Freedom and Operation Iraqi Freedom with posttraumatic stress disorder: a pilot study. *Mil. Med.* 176, 626–630.
44. Wolf, G.K., Stron, T.Q., Kehle, S.M., and Eftekhari, A. (2012). A preliminary examination of prolonged exposure therapy with Iraq and Afghanistan veterans with a diagnosis of posttraumatic stress disorder and mild to moderate traumatic brain injury. *J. Head Trauma Rehabil.* 27, 26–32.
45. Harch, P.G. (2002). The dosage of hyperbaric oxygen in chronic brain injury, in: *The Proceedings of the 2nd International Symposium on Hyperbaric Oxygenation for Cerebral Palsy and the Brain-Injured Child*. J.T. Joiner (ed). Best Publishing Co.: Flagstaff, pps. 31–56.
46. Scorza, K.A., McCarthy, W., Miller, R.S., Carne, W., and Wolf, G. (2013). Hyperbaric oxygen effects on PTSD and mTBI symptoms: a subset analysis. Presented at the Undersea and Hyperbaric Medical Society Annual Meeting, Orlando, FL.
47. Gordillo, G.M., Roy, S., Khanna, S., Schlanger, R., Khandelwal, S., Phillips, G., and Sen, C.K. (2008). Topical oxygen therapy induces vascular endothelial growth factor expression and improves closure of clinically presented chronic wounds. *Clin. Exp. Pharmacol. Physiol.* 35, 957–964.
48. Patel, V., Chivukula, I.V., Roy, S., Khanna, S., He, G., Ojha, N., Mehrotra, A., Dias, L.M., Hunt, T.K., and Sen, C.K. (2005). Oxygen: from the benefits of inducing VEGF expression to managing the risk of hyperbaric stress. *Antioxid. Redox Signal* 7, 1377–1387.
49. Questioning the Hawthorne effect. Light work. Being watched may not affect behaviour, after all. Available at: http://www.economist.com/node/13788427?story_id=13788427. Accessed September 21, 2013.
50. For combat vets, brain injury symptoms can last years. Available at: <http://health.usnews.com/health-news/news/articles/2012/06/20/for-combat-vets-brain-injury-symptoms-can-last-years>. Accessed September 21, 2013.
51. The Veterans Health Administration's treatment of PTSD and traumatic brain injury among recent combat veterans. Available at: <http://www.cbo.gov/sites/default/files/cbofiles/attachments/02-09-PTSD.pdf>. Accessed September 21, 2013.
52. Tunis, S.R., and Pearson, S.D. (2006). Coverage options for promising technologies: Medicare's 'coverage with evidence development.' *Health Aff. (Millwood)* 25, 1218–1230.

Address correspondence to:

Paul G. Harch, MD

Department of Medicine

Louisiana State University School of Medicine

1816 Industrial Boulevard

New Orleans, LA 70058

E-mail: paulharchmd@gmail.com